

X100/301

NATIONAL
QUALIFICATIONS
2010

FRIDAY, 21 MAY
9.00 AM – 10.30 AM

MATHEMATICS
HIGHER
Paper 1
(Non-calculator)

Read carefully

Calculators may NOT be used in this paper.

Section A – Questions 1–20 (40 marks)

Instructions for completion of **Section A** are given on page two.

For this section of the examination you must use an **HB pencil**.

Section B (30 marks)

- 1 Full credit will be given only where the solution contains appropriate working.
- 2 Answers obtained by readings from scale drawings will not receive any credit.



FORMULAE LIST

Circle:

The equation $x^2 + y^2 + 2gx + 2fy + c = 0$ represents a circle centre $(-g, -f)$ and radius $\sqrt{g^2 + f^2 - c}$.

The equation $(x - a)^2 + (y - b)^2 = r^2$ represents a circle centre (a, b) and radius r .

Scalar Product: $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta$, where θ is the angle between $\mathbf{a}$ and $\mathbf{b}$

or $\mathbf{a} \cdot \mathbf{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$ where $\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$.

Trigonometric formulae: $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$= 2\cos^2 A - 1$$

$$= 1 - 2\sin^2 A$$

Table of standard derivatives:

$f(x)$	$f'(x)$
$\sin ax$	$a \cos ax$
$\cos ax$	$-a \sin ax$

Table of standard integrals:

$f(x)$	$\int f(x) dx$
$\sin ax$	$-\frac{1}{a} \cos ax + C$
$\cos ax$	$\frac{1}{a} \sin ax + C$

[Turn over

SECTION A

ALL questions should be attempted.

1. A line L is perpendicular to the line with equation $2x - 3y - 6 = 0$.

What is the gradient of the line L?

A $-\frac{3}{2}$

B $-\frac{1}{2}$

C $\frac{2}{3}$

D 2

2. A sequence is defined by the recurrence relation $u_{n+1} = 2u_n + 3$ and $u_0 = 1$.

What is the value of u_2 ?

A 7

B 10

C 13

D 16

3. Given that $\mathbf{u} = \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} -1 \\ 2 \\ 4 \end{pmatrix}$, find $3\mathbf{u} - 2\mathbf{v}$ in component form.


A $\begin{pmatrix} 4 \\ -1 \\ -5 \end{pmatrix}$

B $\begin{pmatrix} 4 \\ -4 \\ 11 \end{pmatrix}$

C $\begin{pmatrix} 8 \\ -1 \\ 5 \end{pmatrix}$

D $\begin{pmatrix} 8 \\ -4 \\ -5 \end{pmatrix}$

4. The diagram shows the graph with equation of the form $y = a\cos bx$ for $0 \leq x \leq 2\pi$.



What is the equation of this graph?

- A $y = 2\cos 3x$
 - B $y = 2\cos 2x$
 - C $y = 3\cos 2x$
 - D $y = 4\cos 3x$
5. When $x^2 + 8x + 3$ is written in the form $(x + p)^2 + q$, what is the value of q ?
- A -19
 - B -13
 - C -5
 - D 19

[Turn over

6. The roots of the equation $kx^2 - 3x + 2 = 0$ are equal.

What is the value of k ?


- A $-\frac{9}{8}$
- B $-\frac{8}{9}$
- C $\frac{8}{9}$
- D $\frac{9}{8}$

7. A sequence is generated by the recurrence relation $u_{n+1} = \frac{1}{4}u_n + 7$, with $u_0 = -2$.

What is the limit of this sequence as $n \rightarrow \infty$?

- A $\frac{1}{28}$
- B $\frac{28}{5}$
- C $\frac{28}{3}$
- D 28

8. The equation of the circle shown in the diagram is $x^2 + y^2 - 6x - 10y + 9 = 0$.
The x -axis and the line l are parallel tangents to the circle.



What is the equation of line l ?

- A $y = 5$
B $y = 10$
C $y = 18$
D $y = 20$
9. Find $\int (2x^{-4} + \cos 5x) dx$.
A $-\frac{2}{5}x^{-5} - 5\sin 5x + c$
B $-\frac{2}{5}x^{-5} + \frac{1}{5}\sin 5x + c$
C $-\frac{2}{3}x^{-3} + \frac{1}{5}\sin 5x + c$
D $-\frac{2}{3}x^{-3} - 5\sin 5x + c$
10. The vectors $x\mathbf{i} + 5\mathbf{j} + 7\mathbf{k}$ and $-3\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ are perpendicular.
What is the value of x ?
A 0
B 1
C $\frac{4}{3}$
D $\frac{10}{3}$

[Turn over

11. Functions f and g are defined on suitable domains by $f(x) = \cos x$ and $g(x) = x + \frac{\pi}{6}$.

What is the value of $f\left(g\left(\frac{\pi}{6}\right)\right)$?

A $\frac{1}{2} + \frac{\pi}{6}$

B $\frac{\sqrt{3}}{2} + \frac{\pi}{6}$

C $\frac{\sqrt{3}}{2}$

D $\frac{1}{2}$

12. If $f(x) = \frac{1}{\sqrt[5]{x}}$, $x \neq 0$, what is $f'(x)$?

A $-\frac{1}{5}x^{-\frac{6}{5}}$

B $-\frac{1}{5}x^{-\frac{4}{5}}$


C $-\frac{5}{2}x^{-\frac{7}{2}}$

D $-\frac{5}{2}x^{-\frac{3}{2}}$


13. Which of the following diagrams shows a parabola with equation $y = ax^2 + bx + c$, where

- $a > 0$
- $b^2 - 4ac > 0$?


A




B



C




D



[Turn over

14. The diagram shows graphs with equations $y = 14 - x^2$ and $y = 2x^2 + 2$.




Which of the following represents the shaded area?

- A $\int_2^{14} (12 - 3x^2) dx$
- B $\int_2^{14} (3x^2 - 12) dx$
- C $\int_{-2}^2 (12 - 3x^2) dx$
- D $\int_{-2}^2 (3x^2 - 12) dx$
15. The derivative of a function f is given by $f'(x) = x^2 - 9$.
Here are two statements about f :
- (1) f is increasing at $x = 1$;
(2) f is stationary at $x = -3$.

Which of the following is true?

- A Neither statement is correct.
- B Only statement (1) is correct.
- C Only statement (2) is correct.
- D Both statements are correct.

16. The diagram shows the graph with equation $y = k(x - 1)^2(x + t)$.



What are the values of k and t ?

	k	t
A	-2	-5
B	-2	5
C	2	-5
D	2	5

17. If $s(t) = t^2 - 5t + 8$, what is the rate of change of s with respect to t when $t = 3$?


- A -5
- B 1
- C 2
- D 9

18. What is the solution of $x^2 + 4x > 0$, where x is a real number?

- A $-4 < x < 0$
- B $x < -4, x > 0$
- C $0 < x < 4$
- D $x < 0, x > 4$


[Turn over

19. The diagram shows the graph of $y = f(x)$ where f is a logarithmic function.



What is $f(x)$?

- A $f(x) = \log_6(x - 3)$
 - B $f(x) = \log_3(x + 3)$
 - C $f(x) = \log_3(x - 3)$
 - D $f(x) = \log_6(x + 3)$
20. The diagram shows the graph of $y = f(2x) - 3$.



What are the coordinates of the turning point on the graph of $y = f(x)$?

- A $(12, 7)$
- B $(12, 1)$
- C $(3, 7)$
- D $(3, 1)$


[END OF SECTION A]

SECTION B

ALL questions should be attempted.

21. Triangle ABC has vertices A(4, 0), B(−4, 16) and C(18, 20), as shown in the diagram opposite.

Medians AP and CR intersect at the point T(6, 12).



- | | | |
|-----|---------------------------------------|---|
| (a) | Find the equation of median BQ. | 3 |
| (b) | Verify that T lies on BQ. | 1 |
| (c) | Find the ratio in which T divides BQ. | 2 |
-
22. (a) (i) Show that $(x - 1)$ is a factor of $f(x) = 2x^3 + x^2 - 8x + 5$.
- (ii) Hence factorise $f(x)$ fully.
- 5
- (b) Solve $2x^3 + x^2 - 8x + 5 = 0$.
- 1
- (c) The line with equation $y = 2x - 3$ is a tangent to the curve with equation $y = 2x^3 + x^2 - 6x + 2$ at the point G.
- Find the coordinates of G.
- 5
- (d) This tangent meets the curve again at the point H.
- Write down the coordinates of H.
- 1

[Turn over for Question 23 on Page fourteen]

23. (a) Diagram 1 shows a right angled triangle, where the line OA has equation $3x - 2y = 0$.

- (i) Show that $\tan a = \frac{3}{2}$.
(ii) Find the value of $\sin a$.




Diagram 1

4

- (b) A second right angled triangle is added as shown in Diagram 2.

The line OB has equation $3x - 4y = 0$.

Find the values of $\sin b$ and $\cos b$.




Diagram 2

4

- (c) (i) Find the value of $\sin(a - b)$.
(ii) State the value of $\sin(b - a)$.

4

[END OF SECTION B]

[END OF QUESTION PAPER]