



National
Qualifications
2017

2017 Mathematics Paper 2

Higher

Finalised Marking Instructions

© Scottish Qualifications Authority 2017

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is reproduced, SQA should be clearly acknowledged as the source. If it is to be used for any other purpose, written permission must be obtained from permissions@sqa.org.uk.

Where the publication includes materials from sources other than SQA (secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the centre's responsibility to obtain the necessary copyright clearance. SQA's NQ Assessment team may be able to direct you to the secondary sources.

These marking instructions have been prepared by examination teams for use by SQA appointed markers when marking external course assessments. This publication must not be reproduced for commercial or trade purposes.

Question			Generic scheme	Illustrative scheme	Max mark
1.	(a)		<ul style="list-style-type: none"> •¹ find mid-point of BC •² calculate gradient of BC •³ use property of perpendicular lines •⁴ determine equation of line in a simplified form 	<ul style="list-style-type: none"> •¹ $(6, -1)$ •² $-\frac{2}{6}$ •³ 3 •⁴ $y = 3x - 19$ 	4

Notes:

1. •⁴ is only available as a consequence of using a perpendicular gradient **and** a midpoint.
2. The gradient of the perpendicular bisector must appear in simplified form at •³ or •⁴ stage for •³ to be awarded.
3. At •⁴, accept $3x - y - 19 = 0$, $3x - y = 19$ or any other rearrangement of the equation where the constant terms have been simplified.

Commonly Observed Responses:

Question		Generic scheme	Illustrative scheme	Max mark
1.	(b)	<ul style="list-style-type: none"> •⁵ use $m = \tan \theta$ •⁶ determine equation of AB 	<ul style="list-style-type: none"> •⁵ 1 •⁶ $y = x - 3$ 	2

Notes:

4. At •⁶, accept $y - x + 3 = 0$, $y - x = -3$ or any other rearrangement of the equation where the constant terms have been simplified.

Commonly Observed Responses:

Question		Generic scheme	Illustrative scheme	Max mark
1.	(c)	<ul style="list-style-type: none"> •⁷ find x or y coordinate •⁸ find remaining coordinate 	<ul style="list-style-type: none"> •⁷ $x = 8$ or $y = 5$ •⁸ $y = 5$ or $x = 8$ 	2

Notes:

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
2.	(a)		<p>Method 1</p> <ul style="list-style-type: none"> •¹ know to use $x=1$ in synthetic division •² complete division, interpret result and state conclusion 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $\begin{array}{r rrrr} 1 & 2 & -5 & 1 & 2 \\ & & 2 & & \end{array}$ •² $\begin{array}{r rrrr} 1 & 2 & -5 & 1 & 2 \\ & & 2 & -3 & -2 \\ \hline & 2 & -3 & -2 & 0 \end{array}$ <p>Remainder = 0 $\therefore (x-1)$ is a factor</p> 	2
			<p>Method 2</p> <ul style="list-style-type: none"> •¹ know to substitute $x=1$ •² complete evaluation, interpret result and state conclusion 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ $2(1)^3 - 5(1)^2 + (1) + 2$ •² $= 0 \therefore (x-1)$ is a factor 	2
			<p>Method 3</p> <ul style="list-style-type: none"> •¹ start long division and find leading term in quotient •² complete division, interpret result and state conclusion 	<p>Method 3</p> <ul style="list-style-type: none"> •¹ $\begin{array}{r} 2x^2 \\ (x-1) \overline{) 2x^3 - 5x^2 + x + 2} \end{array}$ •² $\begin{array}{r} 2x^2 - 3x - 2 \\ (x-1) \overline{) 2x^3 - 5x^2 + x + 2} \\ \underline{2x^3 - 2x^2} \\ -3x^2 + x \\ \underline{-3x^2 + 3x} \\ -2x + 2 \\ \underline{-2x + 2} \\ 0 \end{array}$ <p>remainder = 0 $\therefore (x-1)$ is a factor</p> 	2

Question	Generic scheme	Illustrative scheme	Max mark
Notes:			
1. Communication at \bullet^2 must be consistent with working at that stage i.e. a candidate's working must arrive legitimately at 0 before \bullet^2 can be awarded. 2. Accept any of the following for \bullet^2 : <ul style="list-style-type: none"> '$f(1) = 0$ so $(x-1)$ is a factor' 'since remainder = 0, it is a factor' the 0 from any method linked to the word 'factor' by e.g. 'so', 'hence', '\therefore', '\rightarrow', '\Rightarrow' 3. Do not accept any of the following for \bullet^2 : <ul style="list-style-type: none"> double underlining the zero or boxing the zero without comment '$x = -1$ is a factor', '$(x+1)$ is a factor', '$(x+1)$ is a root', '$x = 1$ is a root', '$(x-1)$ is a root' '$x = -1$ is a root'. the word 'factor' only with no link 			
Commonly Observed Responses:			

Question	Generic scheme	Illustrative scheme	Max mark
2. (b)	\bullet^3 state quadratic factor \bullet^4 find remaining factors \bullet^5 state solution	\bullet^3 $2x^2 - 3x - 2$ \bullet^4 $(2x+1)$ and $(x-2)$ \bullet^5 $x = -\frac{1}{2}, 1, 2$	3
Notes:			
4. The appearance of " $= 0$ " is not required for \bullet^5 to be awarded. 5. Candidates who identify a different initial factor and subsequent quadratic factor can gain all available marks. 6. \bullet^5 is only available as a result of a valid strategy at \bullet^3 and \bullet^4 . 7. Accept $\left(-\frac{1}{2}, 0\right), (1, 0), (2, 0)$ for \bullet^5 .			
Commonly Observed Responses:			

Question			Generic scheme	Illustrative scheme	Max mark
3.			<ul style="list-style-type: none"> •¹ substitute for y •² express in standard quadratic form •³ factorise •⁴ find x coordinates •⁵ find y coordinates 	<ul style="list-style-type: none"> •¹ $(x-2)^2 + (3x-1)^2 = 25$ or $x^2 - 4x + 4 + (3x)^2 - 2(3x) + 1 = 25$ •² $10x^2 - 10x - 20 = 0$ •³ $10(x-2)(x+1) = 0$ <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <ul style="list-style-type: none"> •⁴ $x = 2$ •⁵ $y = 6$ </div> <div style="text-align: center;"> <ul style="list-style-type: none"> •⁵ $x = -1$ $y = -3$ </div> </div> 	5

Notes:

1. At •³ the quadratic must lead to two distinct real roots for •⁴ and •⁵ to be available.
2. •² is only available if '= 0' appears at •² or •³ stage.
3. If a candidate arrives at an equation which is not a quadratic at •² stage, then •³, •⁴ and •⁵ are not available
4. At •³ do not penalise candidates who fail to extract the common factor or who have divided the quadratic equation by 10.
5. •³ is available for substituting correctly into the quadratic formula.
6. •⁴ and •⁵ may be marked either horizontally or vertically.
7. For candidates who identify **both** solutions by inspection, full marks may be awarded provided they justify that their points lie on **both** the line and the circle. Candidates who identify **both** solutions, but justify only one gain 2 out of 5.

Commonly Observed Responses:

Candidate A

$$(x-2)^2 + (3x-1)^2 = 25 \quad \bullet^1 \checkmark$$

$$10x^2 - 10x = 20 \quad \bullet^2 \times$$

$$10x(x-1) = 20 \quad \bullet^3 \boxed{\checkmark 2}$$

$$x = 2 \quad x = 3 \quad \bullet^4 \times$$

$$y = 6 \quad y = 9 \quad \bullet^5 \boxed{\checkmark 2}$$

Candidate B

Candidates who substitute into the circle equation only

$$\bullet^1 \checkmark$$

$$\bullet^2 \checkmark$$

$$\bullet^3 \checkmark$$

$$\bullet^4 \checkmark$$

Sub $x = 2$

$$y^2 - 2y - 24 = 0$$

$$(y-6)(y+4) = 0$$

$$y = 6 \text{ or } y = -4$$

Sub $x = -1$

$$y^2 - 2y - 15 = 0$$

$$(y+3)(y-5) = 0$$

$$y = -3 \text{ or } y = 5$$

(2, 6) (-1, -3) $\bullet^5 \times$

Question			Generic scheme	Illustrative scheme	Max mark
4.	(a)		Method 1 <ul style="list-style-type: none"> •¹ identify common factor •² complete the square •³ process for c and write in required form 	Method 1 <ul style="list-style-type: none"> •¹ $3(x^2 + 8x \dots \dots)$ stated or implied by •² •² $3(x+4)^2 \dots \dots$ •³ $3(x+4)^2 + 2$ 	3
			Method 2 <ul style="list-style-type: none"> •¹ expand completed square form •² equate coefficients •³ process for b and c and write in required form 	Method 2 <ul style="list-style-type: none"> •¹ $ax^2 + 2abx + ab^2 + c$ •² $a = 3, 2ab = 24, ab^2 + c = 50$ •³ $3(x+4)^2 + 2$ 	3

Notes:

1. $3(x+4)^2 + 2$ with no working gains •¹ and •² only; however, see Candidate G.
2. •³ is only available for a calculation involving both multiplication and subtraction of integers.

Commonly Observed Responses:

Candidate A $3\left(x^2 + 8x + \frac{50}{3}\right)$ $3\left(x^2 + 8x + 16 - 16 + \frac{50}{3}\right)$ <ul style="list-style-type: none"> •¹ ✓ •² ^ further working is required 	Candidate B $3x^2 + 24x + 50 = 3(x+8)^2 - 64 + 50$ $= 3(x+8)^2 - 14$ <ul style="list-style-type: none"> •¹ ✗ •² ✗ •³ ✓2
Candidate C $ax^2 + 2abx + ab^2 + c$ $a = 3, 2ab = 24, b^2 + c = 50$ $a = 3, b = 4, c = 34$ $3(x+4)^2 + 34$ <ul style="list-style-type: none"> •¹ ✓ •² ✗ •³ ✓1 	Candidate D $3((x^2 + 24x) + 50)$ $3((x+12)^2 - 144) + 50$ $3(x+12)^2 - 382$ <ul style="list-style-type: none"> •¹ ✗ •² ✓1 •³ ✓1

Question	Generic scheme	Illustrative scheme	Max mark
Candidate E $a(x+b)^2 + c = ax^2 + 2abx + ab^2 + c$ • ¹ ✓ $a = 3, 2ab = 24, ab^2 + c = 50$ • ² ✓ $b = 4, c = 2$ • ³ ✓ <div>•³ is awarded as all working relates to completed square form</div>	Candidate F $ax^2 + 2abx + ab^2 + c$ • ¹ ✓ $a = 3, 2ab = 24, ab^2 + c = 50$ • ² ✓ $b = 4, c = 2$ • ³ ✗ <div>•³ is lost as no reference is made to completed square form</div>		
Candidate G $3(x+4)^2 + 2$ Check: $3(x^2 + 8x + 16) + 2$ $= 3x^2 + 24x + 48 + 2$ $= 3x^2 + 24x + 50$ Award 3/3	Candidate H $3x^2 + 24x + 50$ $= 3(x+4)^2 - 16 + 50$ • ¹ ✓ • ² ✓ $= 3(x+4)^2 + 34$ • ³ ✗		

Question	Generic scheme	Illustrative scheme	Max mark
4. (b)	• ⁴ differentiate two terms • ⁵ complete differentiation	• ⁴ $3x^2 + 24x....$ • ⁵ $.... + 50$	2
Notes: 3. • ⁴ is awarded for any two of the following three terms: $3x^2$, $+24x$, $+50$			
Commonly Observed Responses:			

Question			Generic scheme	Illustrative scheme	Max mark
4.	(c)		<p>Method 1</p> <ul style="list-style-type: none"> •⁶ link with (a) and identify sign of $(x+4)^2$ •⁷ communicate reason <p>Method 2</p> <ul style="list-style-type: none"> •⁶ identify minimum value of $f'(x)$ •⁷ communicate reason 	<p>Method 1</p> <ul style="list-style-type: none"> •⁶ $f'(x) = 3(x+4)^2 + 2$ and $(x+4)^2 \geq 0 \forall x$ •⁷ $\therefore 3(x+4)^2 + 2 > 0 \Rightarrow$ always strictly increasing <p>Method 2</p> <ul style="list-style-type: none"> •⁶ eg minimum value = 2 or annotated sketch •⁷ $2 > 0 \therefore (f'(x) > 0) \Rightarrow$ always strictly increasing 	2

Notes:

- Do not penalise $(x+4)^2 > 0$ or the omission of $f'(x)$ at •⁶ in Method 1.
- Responses in part (c) must be consistent with working in parts (a) and (b) for •⁶ and •⁷ to be available.
- Where erroneous working leads to a candidate considering a function which is not always strictly increasing, only •⁶ is available.
- At •⁶ communication should be explicitly in terms of the given function. Do not accept statements such as “(something)² ≥ 0 ”, “something squared ≥ 0 ”. However, •⁷ is still available.

Commonly Observed Responses:

<p>Candidate I</p> $f'(x) = 3(x+4)^2 + 2$ $3(x+4)^2 + 2 > 0 \Rightarrow \text{strictly increasing.}$ <p>Award 1 out of 2</p>	<p>Candidate J</p> <p>Since $3x^2 + 24x + 50 = 3(x+4)^2 + \frac{166}{50}$</p> <p>and $(x+4)^2$ is > 0 for all x then</p> $3(x+4)^2 + \frac{166}{50} > 0 \text{ for all } x.$ <p>Hence the curve is strictly increasing for all values of x. •⁶ ✓ •⁷ ✓1</p>
---	---

Question			Generic scheme	Illustrative scheme	Max mark
5.	(a)		• ¹ identify pathway • ² state \overrightarrow{PQ}	• ¹ $\overrightarrow{PR} + \overrightarrow{RQ}$ stated or implied by • ² • ² $-3\mathbf{i} - 4\mathbf{j} + 5\mathbf{k}$	2

Notes:

- Award •¹ $(9\mathbf{i} + 5\mathbf{j} + 2\mathbf{k}) + (-12\mathbf{i} - 9\mathbf{j} + 3\mathbf{k})$.
- Candidates who choose to work with column vectors and leave their answer in the form $\begin{pmatrix} -3 \\ -4 \\ 5 \end{pmatrix}$ cannot gain •².
- ² is not available for simply adding or subtracting vectors within an invalid strategy.
- Where candidates choose specific points consistent with the given vectors, only •¹ and •⁴ are available. However, should the statement 'without loss of generality' precede the selected points then marks •¹, •², •³ and •⁴ are all available.

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
5.	(b)		• ³ interpret ratio • ⁴ identify pathway and demonstrate result	• ³ $\frac{2}{3}$ or $\frac{1}{3}$ • ⁴ $\overrightarrow{PR} + \frac{2}{3}\overrightarrow{RQ}$ or $\overrightarrow{PQ} + \frac{1}{3}\overrightarrow{QR}$ leading to $\mathbf{i} - \mathbf{j} + 4\mathbf{k}$	2

Notes:

- This is a 'show that' question. Candidates who choose to work with column vectors must write their final answer in the required form to gain •⁴. $\begin{pmatrix} 1 \\ -1 \\ 4 \end{pmatrix}$ does not gain •⁴.
- Beware of candidates who fudge their working between •³ and •⁴.

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
<p>Candidate A - legitimate use of the section formula</p> $\overrightarrow{PS} = \frac{n\overrightarrow{PQ} + m\overrightarrow{PR}}{m+n}$ $\overrightarrow{PS} = \frac{2\overrightarrow{PQ} + \overrightarrow{PR}}{3} \quad \bullet^3 \checkmark$ $\overrightarrow{PS} = \frac{2\begin{pmatrix} -3 \\ -4 \\ 5 \end{pmatrix} + \begin{pmatrix} 9 \\ 5 \\ 2 \end{pmatrix}}{3} + \frac{\begin{pmatrix} 9 \\ 5 \\ 2 \end{pmatrix}}{3}$ $= \begin{pmatrix} -2 \\ -8/3 \\ 10/3 \end{pmatrix} + \begin{pmatrix} 3 \\ 5/3 \\ 2/3 \end{pmatrix}$ $= \begin{pmatrix} 1 \\ -1 \\ 4 \end{pmatrix}$ $\overrightarrow{PS} = \mathbf{i} - \mathbf{j} + 4\mathbf{k} \quad \bullet^4 \checkmark$		<p>Candidate B - BEWARE - treating P as the origin</p> $2\overrightarrow{QS} = \overrightarrow{SR}$ $3\mathbf{s} = 2\mathbf{q} + \mathbf{r} \quad \bullet^3 \checkmark$ $3\mathbf{s} = 2\begin{pmatrix} -3 \\ -4 \\ 5 \end{pmatrix} + \begin{pmatrix} 9 \\ 5 \\ 2 \end{pmatrix}$ $\mathbf{s} = \mathbf{i} - \mathbf{j} + 4\mathbf{k} \quad \bullet^4 \times$	

Question			Generic scheme	Illustrative scheme	Max mark
5.	(c)		<p>Method 1</p> <ul style="list-style-type: none"> •⁵ evaluate $\overrightarrow{PQ} \cdot \overrightarrow{PS}$ •⁶ evaluate \overrightarrow{PQ} •⁷ evaluate \overrightarrow{PS} •⁸ use scalar product •⁹ calculate angle 	<p>Method 1</p> <ul style="list-style-type: none"> •⁵ $\overrightarrow{PQ} \cdot \overrightarrow{PS} = 21$ •⁶ $\overrightarrow{PQ} = \sqrt{50}$ •⁷ $\overrightarrow{PS} = \sqrt{18}$ •⁸ $\cos QPS = \frac{21}{\sqrt{50} \times \sqrt{18}}$ •⁹ 45.6° or 0.795 radians 	5
			<p>Method 2</p> <ul style="list-style-type: none"> •⁵ evaluate \overrightarrow{QS} •⁶ evaluate \overrightarrow{PQ} •⁷ evaluate \overrightarrow{PS} •⁸ use cosine rule •⁹ calculate angle 	<p>Method 2</p> <ul style="list-style-type: none"> •⁵ $\overrightarrow{QS} = \sqrt{26}$ •⁶ $\overrightarrow{PQ} = \sqrt{50}$ •⁷ $\overrightarrow{PS} = \sqrt{18}$ •⁸ $\cos QPS = \frac{(\sqrt{50})^2 + (\sqrt{18})^2 - (\sqrt{26})^2}{2 \times \sqrt{50} \times \sqrt{18}}$ •⁹ 45.6° or 0.795 radians 	5

Notes:

- For candidates who use \overrightarrow{PS} not equal to $\mathbf{i} - \mathbf{j} + 4\mathbf{k}$ •⁵ is not available in Method 1 or •⁷ in Method 2.
 - Do not penalise candidates who treat negative signs with a lack of rigour when calculating a magnitude. However, $\sqrt{1^2 - 1^2 + 4^2}$ leading to $\sqrt{16}$ indicates an invalid method for calculating the magnitude. No mark can be awarded for any magnitude arrived at using an invalid method.
 - ⁸ is not available to candidates who simply state the formula $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}| |\mathbf{b}|}$.
- However, $\cos \theta = \frac{\overrightarrow{PQ} \cdot \overrightarrow{PS}}{|\overrightarrow{PQ}| \times |\overrightarrow{PS}|}$ or $\cos \theta = \frac{21}{\sqrt{50} \times \sqrt{18}}$ is acceptable. Similarly for Method 2.
- Accept answers which round to 46° or 0.8 radians.
 - Do not penalise the omission or incorrect use of units.
 - ⁹ is only available as a result of using a valid strategy.
 - ⁹ is only available for a single angle.
 - For a correct answer with no working award 0/5.

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
Candidate C - Calculating wrong angle		Candidate D- Calculating wrong angle	
$\overrightarrow{QP} \cdot \overrightarrow{QS} = 29$ • ⁵ ✗ $ \overrightarrow{QP} = \sqrt{50}$ • ⁶ ✓1 $ \overrightarrow{QS} = \sqrt{26}$ • ⁷ ✓1 $\cos \hat{PQS} = \frac{29}{\sqrt{50} \times \sqrt{26}}$ • ⁸ ✓1 $\hat{PQS} = 36.5$ • ⁹ ✗ strategy incomplete		$\overrightarrow{PS} \cdot \overrightarrow{QP} = -21$ • ⁵ ✗ $ \overrightarrow{QP} = \sqrt{50}$ • ⁶ ✓ $ \overrightarrow{PS} = \sqrt{18}$ • ⁷ ✓ $\cos \theta = \frac{-21}{\sqrt{50} \times \sqrt{18}}$ • ⁸ ✓1 $\theta = 134.4$ • ⁹ ✗ strategy incomplete	
For candidates who continue, and use the angle found to evaluate the required angle, then all marks are available.		For candidates who continue, and use the angle found to evaluate the required angle, then all marks are available.	
Candidate E From (a) $\overrightarrow{PQ} = -21\mathbf{i} - 14\mathbf{j} + \mathbf{k}$		Candidate F From (a) $\overrightarrow{PQ} = 21\mathbf{i} + 14\mathbf{j} - \mathbf{k}$	
$\overrightarrow{PQ} \cdot \overrightarrow{PS} = -3$ • ⁵ ✓1 $ \overrightarrow{PQ} = \sqrt{638}$ • ⁶ ✓1 $ \overrightarrow{PS} = \sqrt{18}$ • ⁷ ✓ $\cos \hat{QPS} = \frac{-3}{\sqrt{638} \times \sqrt{18}}$ • ⁸ ✓1 $\hat{QPS} = 91.6$ • ⁹ ✓1		$\overrightarrow{PQ} \cdot \overrightarrow{PS} = 3$ • ⁵ ✓1 $ \overrightarrow{PQ} = \sqrt{638}$ • ⁶ ✓1 $ \overrightarrow{PS} = \sqrt{18}$ • ⁷ ✓ $\cos \hat{QPS} = \frac{3}{\sqrt{638} \times \sqrt{18}}$ • ⁸ ✓1 $\hat{QPS} = 88.4$ • ⁹ ✓1	
Candidate G From (b) $\overrightarrow{PS} = -4\mathbf{i} - 3\mathbf{j} + \mathbf{k}$			
$\overrightarrow{PQ} \cdot \overrightarrow{PS} = 3$ • ⁵ ✗ $ \overrightarrow{PQ} = \sqrt{50}$ • ⁶ ✓ $ \overrightarrow{PS} = \sqrt{26}$ • ⁷ ✓1 $\cos \hat{QPS} = \frac{3}{\sqrt{50} \times \sqrt{26}}$ • ⁸ ✓1 $\hat{QPS} = 85.2$ • ⁹ ✓1			

Question			Generic scheme	Illustrative scheme	Max mark
6.			<ul style="list-style-type: none"> •¹ substitute appropriate double angle formula •² express in standard quadratic form •³ factorise •⁴ solve for $\sin x^\circ$ •⁵ solve for x 	<ul style="list-style-type: none"> •¹ $5\sin x - 4 = 2(1 - 2\sin^2 x)$ •² $4\sin^2 x + 5\sin x - 6 = 0$ •³ $(4\sin x - 3)(\sin x + 2)$ •⁴ $\sin x = \frac{3}{4},$ •⁵ $\sin x = -2$ •⁵ $x = 0.848, 2.29, \cancel{\sin x = -2}$ 	5

Notes:

1. •¹ is not available for simply stating $\cos 2x = 1 - 2\sin^2 x$ with no further working.
2. In the event of $\cos^2 x^\circ - \sin^2 x^\circ$ or $2\cos^2 x^\circ - 1$ being substituted for $\cos 2x$, •¹ cannot be awarded until the equation reduces to a quadratic in $\sin x^\circ$.
3. Substituting $1 - 2\sin^2 A$ or $1 - 2\sin^2 \alpha$ for $\cos 2x$ at •¹ stage should be treated as bad form provided the equation is written in terms of x at •² stage. Otherwise, •¹ is not available.
4. ' $= 0$ ' must appear by •³ stage for •² to be awarded. However, for candidates using the quadratic formula to solve the equation, ' $= 0$ ' must appear at •² stage for •² to be awarded.
5. $5\sin x + 4\sin^2 x - 6 = 0$ does not gain •² unless •³ is awarded.
6. $\sin x = \frac{-5 \pm \sqrt{121}}{8}$ gains •³.
7. Candidates may express the equation obtained at •² in the form $4s^2 + 5s - 6 = 0$ or $4x^2 + 5x - 6 = 0$. In these cases, award •³ for $(4s - 3)(s + 2) = 0$ or $(4x - 3)(x + 2) = 0$. However, •⁴ is only available if $\sin x$ appears explicitly at this stage.
8. •⁴ and •⁵ are only available as a consequence of solving a quadratic equation.
9. •³, •⁴ and •⁵ are not available for any attempt to solve a quadratic equation written in the form $ax^2 + bx = c$.
10. •⁵ is not available to candidates who work in degrees and do not convert their solutions into radian measure.
11. Accept answers which round to 0.85 and 2.3 at •⁵ eg $\frac{49\pi}{180}, \frac{131\pi}{180}$.
12. Answers written as decimals should be rounded to no fewer than 2 significant figures.
13. Do not penalise additional solutions at •⁵.

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
Candidate A		Candidate B	
$\bullet^1 \checkmark \bullet^2 \checkmark$ $(4s-3)(s+2) = 0$ $s = \frac{3}{4}, s = -2$ $x = 0.848, 2.29$	$\bullet^3 \checkmark$ $\bullet^4 \times$ $\bullet^5 \checkmark$	$\bullet^1 \checkmark$ $4\sin^2 x + 5\sin x - 6 = 0$ $9\sin x - 6 = 0$ $\sin x = \frac{2}{3}$ $x = 0.730, 2.41$	$\bullet^2 \checkmark$ $\bullet^3 \times$ $\bullet^4 \boxed{\checkmark 2}$ $\bullet^5 \boxed{\checkmark 2}$
Candidate C		Candidate D	
$5\sin x - 4 = 2(1 - 2\sin^2 x)$ $4\sin^2 x + 5\sin x = 6$ $\sin x(4\sin x + 5) = 6$ $\sin x = 6, 4\sin x + 5 = 6$ no solution, $\sin x = \frac{1}{4}$ $x = 0.253, 2.89$	$\bullet^1 \checkmark$ $\bullet^2 \boxed{\checkmark 2}$ $\bullet^3 \boxed{\checkmark 2}$ $\bullet^4 \times$ $\bullet^5 \times$	$5\sin x - 4 = 2(1 - 2\sin^2 x)$ $4\sin^2 x + 5\sin x - 6 = 0$ $4\sin^2 x + 5\sin x = 6$ $\sin x(4\sin x + 5) = 6$ $\sin x = 6, 4\sin x + 5 = 6$ no solution, $\sin x = \frac{1}{4}$ $x = 0.253, 2.89$	$\bullet^1 \checkmark$ $\bullet^2 \checkmark$ $\bullet^3 \boxed{\checkmark 2}$ $\bullet^4 \times$ $\bullet^5 \times$
Candidate E - reading $\cos 2x$ as $\cos^2 x$			
$5\sin x - 4 = 2\cos^2 x$ $5\sin x - 4 = 2(1 - \sin^2 x)$ $2\sin^2 x + 5\sin x - 6 = 0$ $\sin x = \frac{-5 \pm \sqrt{73}}{4}$ $\sin x = 0.886, \sin x = -3.386$ $x = 1.08, 2.05$	$\bullet^1 \times$ $\bullet^2 \boxed{\checkmark 1}$ $\bullet^3 \boxed{\checkmark 1}$ $\bullet^4 \boxed{\checkmark 1}$ $\bullet^5 \boxed{\checkmark 1}$		

Question			Generic scheme	Illustrative scheme	Max mark
7.	(a)		<ul style="list-style-type: none"> •¹ write in differentiable form •² differentiate one term •³ complete differentiation and equate to zero •⁴ solve for x 	<ul style="list-style-type: none"> •¹ $\dots - 2x^{\frac{3}{2}}$ stated or implied •² $\frac{dy}{dx} = 6\dots$ or $\frac{dy}{dx} = \dots - 3x^{\frac{1}{2}}\dots$ •³ $\dots - 3x^{\frac{1}{2}} = 0$ or $6\dots = 0$ •⁴ $x = 4$ 	4

Notes:

1. For candidates who do not differentiate a term involving a fractional index, either •² or •³ is available but not both.
2. •⁴ is available only as a consequence of solving an equation involving a fractional power of x .
3. For candidates who integrate one or other of the terms •⁴ is unavailable.

Commonly Observed Responses:

Candidate A - differentiating incorrectly

$$y = 6x - 2x^{\frac{3}{2}} \quad \bullet^1 \checkmark$$

$$\frac{dy}{dx} = 6 - 3x^{\frac{5}{2}} \quad \bullet^2 \checkmark$$

$$6 - 3x^{\frac{5}{2}} = 0 \quad \bullet^3 \times$$

$$x = 1.32 \quad \bullet^4 \boxed{\checkmark 1}$$

Candidate B - integrating the second term

$$y = 6x - 2x^{\frac{3}{2}} \quad \bullet^1 \checkmark$$

$$\frac{dy}{dx} = 6 - \frac{4}{5}x^{\frac{5}{2}} \quad \bullet^2 \checkmark$$

$$6 - \frac{4}{5}x^{\frac{5}{2}} = 0 \quad \bullet^3 \times$$

$$x = 2.24 \quad \bullet^4 \times$$

Question			Generic scheme	Illustrative scheme	Max mark
7.	(b)		<ul style="list-style-type: none"> •⁵ evaluate y at stationary point •⁶ consider value of y at end points •⁷ state greatest and least values 	<ul style="list-style-type: none"> •⁵ 8 •⁶ 4 and 0 •⁷ greatest 8, least 0 stated explicitly 	3

Notes:

- The only valid approach to finding the stationary point is via differentiation. A numerical approach can only gain •⁶.
- ⁷ is not available to candidates who do not consider both end points.
- Vertical marking is not applicable to •⁶ and •⁷.
- Ignore any nature table which may appear in a candidate's solution; however, the appearance of (4,8) at a nature table is sufficient for •⁵.
- Greatest (4,8); least (9,0) does not gain •⁷.
- ⁵ and •⁷ are not available for evaluating y at a value of x , obtained at •⁴ stage, which lies outwith the interval $1 \leq x \leq 9$.
- For candidates who only evaluate the derivative, •⁵, •⁶ and •⁷ are not available.

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
8.	(a)		<ul style="list-style-type: none"> •¹ find expression for u_1 •² find expression for u_2 and express in the correct form 	<ul style="list-style-type: none"> •¹ $5k - 20$ •² $u_2 = k(5k - 20) - 20$ leading to $u_2 = 5k^2 - 20k - 20$ 	2

Notes:

Commonly Observed Responses:

Question			Generic scheme	Illustrative scheme	Max mark
8.	(b)		<ul style="list-style-type: none"> •³ interpret information •⁴ express inequality in standard quadratic form •⁵ determine zeros of quadratic expression •⁶ state range with justification 	<ul style="list-style-type: none"> •³ $5k^2 - 20k - 20 < 5$ •⁴ $5k^2 - 20k - 25 < 0$ •⁵ $-1, 5$ •⁶ $-1 < k < 5$ with eg sketch or table of signs 	4
Notes:					
<p>1. Candidates who work with an equation from the outset lose •³ and •⁴. However, •⁵ and •⁶ are still available.</p> <p>2. At •⁵ do not penalise candidates who fail to extract the common factor or who have divided the quadratic inequation by 5.</p> <p>3. •⁴ and •⁵ are only available to candidates who arrive at a quadratic expression at •³.</p> <p>4. At •⁶ accept “$k > -1$ and $k < 5$” or “$k > -1, k < 5$” together with the required justification.</p> <p>5. For a trial and error approach award 0/4.</p>					
Commonly Observed Responses:					

Question			Generic scheme	Illustrative scheme	Max mark
9.			<p>Method 1</p> <ul style="list-style-type: none"> •¹ state linear equation •² introduce logs •³ use laws of logs •⁴ use laws of logs •⁵ state k and n 	<p>Method 1</p> <ul style="list-style-type: none"> •¹ $\log_2 y = \frac{1}{4}\log_2 x + 3$ •² $\log_2 y = \frac{1}{4}\log_2 x + 3\log_2 2$ •³ $\log_2 y = \log_2 x^{\frac{1}{4}} + \log_2 2^3$ •⁴ $\log_2 y = \log_2 2^3 x^{\frac{1}{4}}$ •⁵ $k = 8, n = \frac{1}{4}$ or $y = 8x^{\frac{1}{4}}$ 	5
			<p>Method 2</p> <ul style="list-style-type: none"> •¹ state linear equation •² use laws of logs •³ use laws of logs •⁴ use laws of logs •⁵ state k and n 	<p>Method 2</p> <ul style="list-style-type: none"> •¹ $\log_2 y = \frac{1}{4}\log_2 x + 3$ •² $\log_2 y = \log_2 x^{\frac{1}{4}} + 3$ •³ $\log_2 \frac{y}{x^{\frac{1}{4}}} = 3$ •⁴ $\frac{y}{x^{\frac{1}{4}}} = 2^3$ •⁵ $k = 8, n = \frac{1}{4}$ or $y = 8x^{\frac{1}{4}}$ 	5

Question			Generic Scheme	Illustrative Scheme	Max Mark
			<p>Method 3</p> <ul style="list-style-type: none"> •¹ introduce logs to $y = kx^n$ •² use laws of logs •³ interpret intercept •⁴ use laws of logs •⁵ interpret gradient 	<p>Method 3</p> <p>The equations at •¹, •² and •³ must be stated explicitly.</p> <ul style="list-style-type: none"> •¹ $\log_2 y = \log_2 kx^n$ •² $\log_2 y = n \log_2 x + \log_2 k$ •³ $\log_2 k = 3$ •⁴ $k = 8$ •⁵ $n = \frac{1}{4}$ 	5
			<p>Method 4</p> <ul style="list-style-type: none"> •¹ interpret point on log graph •² convert from log to exp. form •³ interpret point and convert •⁴ substitute into $y = kx^n$ and evaluate k •⁵ substitute other point into $y = kx^n$ and evaluate n 	<p>Method 4</p> <ul style="list-style-type: none"> •¹ $\log_2 x = -12$ and $\log_2 y = 0$ •² $x = 2^{-12}$ and $y = 2^0$ •³ $\log_2 x = 0, \log_2 y = 3$ $x = 1, y = 2^3$ •⁴ $2^3 = k \times 1^n \Rightarrow k = 8$ •⁵ $2^0 = 2^3 \times 2^{-12n}$ $\Rightarrow 3 - 12n = 0$ $\Rightarrow n = \frac{1}{4}$ 	5
Notes: <ol style="list-style-type: none"> Markers must not pick and choose between methods. Identify the method which best matches the candidates approach. Treat the omission of base 2 as bad form at •¹ and •³ in Method 1, at •¹ and •² for Method 2 and Method 3, and at •¹ in Method 4. '$m = \frac{1}{4}$', or 'gradient = $\frac{1}{4}$', does not gain •⁵ in Method 3. Accept 8 in lieu of 2^3 throughout. In Method 4 candidates may use (0,3) for •¹ and •² followed by (-12,0) for •³. 					

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
<p>Candidate A</p> <p>With no working. Method 3:</p> $k = 8$ $n = \frac{1}{4}$ <p>Award 2/5</p>	<p>•⁴ ✓</p> <p>•⁵ ✓</p>	<p>Candidate B</p> <p>With no working. Method 3:</p> $n = 8$ $k = \frac{1}{4}$ <p>Award 0/5</p>	<p>•⁴ ✗</p> <p>•⁵ ✗</p>
<p>Candidate C</p> <p>Method 3:</p> $\log_2 k = 3$ $k = 8$ $n = \frac{1}{4}$ <p>Award 3/5</p>	<p>•³ ✓</p> <p>•⁴ ✓</p> <p>•⁵ ✓</p>	<p>Candidate D</p> <p>Method 2:</p> $\log_2 y = \frac{1}{4} \log_2 x + 3$ $\log_2 y = \log_2 x^{\frac{1}{4}} + 3$ $y = x^{\frac{1}{4}} + 3$ $k = 1, n = \frac{1}{4}$ <p>Award 2/5</p>	<p>•¹ ✓</p> <p>•² ✓</p> <p>•³ ✗ •⁴ ✗</p> <p>•⁵ ✗</p>
<p>Candidate E</p> <p>Method 2:</p> $y = \frac{1}{4}x + 3$ <p>~~~~~</p> $\log_2 y = \frac{1}{4} \log_2 x + 3$ $\log_2 y = \log_2 x^{\frac{1}{4}} + 3$ $\frac{y}{x^{\frac{1}{4}}} = 3$ $y = 3x^{\frac{1}{4}}$ <p>Award 3/5</p>	<p>•¹ ✓</p> <p>•² ✓</p> <p>•³ ^ •⁴ ✗</p> <p>•⁵ ✓1</p>		

Question			Generic scheme	Illustrative scheme	Max mark
10.	(a)		Method 1 <ul style="list-style-type: none"> •¹ calculate m_{AB} •² calculate m_{BC} •³ interpret result and state conclusion 	Method 1 <ul style="list-style-type: none"> •¹ $m_{AB} = \frac{3}{9} = \frac{1}{3}$ see Note 1 •² $m_{BC} = \frac{5}{15} = \frac{1}{3}$ •³ ... \Rightarrow AB and BC are parallel (common direction), B is a common point, hence A, B and C are collinear. 	3
			Method 2 <ul style="list-style-type: none"> •¹ calculate an appropriate vector e.g. \overline{AB} •² calculate a second vector e.g. \overline{BC} and compare •³ interpret result and state conclusion 	Method 2 <ul style="list-style-type: none"> •¹ $\overline{AB} = \begin{pmatrix} 9 \\ 3 \end{pmatrix}$ see Note 1 •² $\overline{BC} = \begin{pmatrix} 15 \\ 5 \end{pmatrix} \therefore \overline{AB} = \frac{3}{5}\overline{BC}$ •³ ... \Rightarrow AB and BC are parallel (common direction), B is a common point, hence A, B and C are collinear. 	3
			Method 3 <ul style="list-style-type: none"> •¹ calculate m_{AB} •² find equation of line and substitute point •³ communication 	Method 3 <ul style="list-style-type: none"> •¹ $m_{AB} = \frac{3}{9} = \frac{1}{3}$ •² eg, $y - 1 = \frac{1}{3}(x - 2)$ leading to $6 - 1 = \frac{1}{3}(17 - 2)$ •³ since C lies on line A, B and C are collinear 	
Notes: <ol style="list-style-type: none"> At •¹ and •² stage, candidates may calculate the gradients/vectors using any pair of points. •³ can only be awarded if a candidate has stated “parallel”, “common point” and “collinear”. Candidates who state “points A, B and C are parallel” or “m_{AB} and m_{BC} are parallel” do not gain •³. 					

Question	Generic scheme	Illustrative scheme	Max mark
Commonly Observed Responses:			
<p>Candidate A</p> $m_{AB} = \frac{3}{9} = \frac{1}{3}$ $m_{BC} = \frac{5}{15}$ <p>\Rightarrow AB and BC are parallel , B is a common point, hence A, B and C are collinear.</p>	<p>Candidate B</p> $\begin{pmatrix} 9 \\ 3 \end{pmatrix}$ $\begin{pmatrix} 15 \\ 5 \end{pmatrix} \therefore \overline{AB} = \frac{5}{3} \overline{BC}$ <p>\Rightarrow AB and BC are parallel , B is a common point, hence A, B and C are collinear.</p>	<p>Candidate C</p> $\overline{AB} = \begin{pmatrix} 9 \\ 3 \end{pmatrix}$ $\overline{BC} = \begin{pmatrix} 15 \\ 5 \end{pmatrix} = 5 \begin{pmatrix} 3 \\ 1 \end{pmatrix} \text{ and}$ $\begin{pmatrix} 9 \\ 3 \end{pmatrix} = 3 \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ <p>$\therefore \overline{AB} = \frac{5}{3} \overline{BC}$ ignore working subsequent to correct statement at \bullet^2. \Rightarrow AB and BC are parallel , B is a common point, hence A, B and C are collinear.</p>	<p>$\bullet^1 \checkmark$</p> <p>$\bullet^2 \wedge$</p> <p>$\bullet^3 \boxed{\checkmark 1}$</p> <p>$\bullet^1 \checkmark$</p> <p>$\bullet^2 \times$</p> <p>$\bullet^3 \boxed{\checkmark 1}$</p> <p>$\bullet^1 \checkmark$</p> <p>$\bullet^2 \checkmark$</p> <p>$\bullet^3 \checkmark$</p>

Question			Generic scheme	Illustrative scheme	Max mark
11.	(a)		Method 1 <ul style="list-style-type: none"> •¹ substitute for $\sin 2x$ •² simplify and factorise •³ substitute for $1 - \cos^2 x$ and simplify 	Method 1 <ul style="list-style-type: none"> •¹ $\frac{2 \sin x \cos x}{2 \cos x} - \sin x \cos^2 x$ stated explicitly as above or in a simplified form of the above •² $\sin x(1 - \cos^2 x)$ •³ $\sin x \times \sin^2 x$ leading to $\sin^3 x$ 	3
			Method 2 <ul style="list-style-type: none"> •¹ substitute for $\sin 2x$ •² simplify and substitute for $\cos^2 x$ •³ expand and simplify 	Method 2 <ul style="list-style-type: none"> •¹ $\frac{2 \sin x \cos x}{2 \cos x} - \sin x \cos^2 x$ stated explicitly as above or in a simplified form of the above •² $\sin x - \sin x(1 - \sin^2 x)$ •³ $\sin x - \sin x + \sin^3 x$ leading to $\sin^3 x$ 	3

Notes:

- ¹ is not available to candidates who simply quote $\sin 2x = 2 \sin x \cos x$ without substituting into the expression given on the LHS. See Candidate B
- In method 2 where candidates attempt •¹ and •² in the same line of working •¹ may still be awarded if there is an error at •².
- ³ is not available to candidates who work throughout with A in place of x .
- Treat multiple attempts which are not scored out as different strategies, and apply General Marking Principle (r).
- On the appearance of $\text{LHS} = 0$, the first available mark is lost; however, any further marks are still available.

Commonly Observed Responses:

Candidate A

$$\frac{2 \sin x \cos x}{2 \cos x} - \sin x \cos^2 x = \sin^3 x \quad \bullet^1 \checkmark$$

$$\sin x - \sin x \cos^2 x = \sin^3 x \quad \bullet^2 \wedge$$

$$1 - \cos^2 x = \sin^2 x \quad \bullet^3 \times$$

$$\sin^2 x = \sin^2 x$$

In proving the identity, candidates must work with both sides independently ie in each line of working the LHS must be equivalent to the line above.

Candidate B

$$\text{LHS} = \frac{\sin 2x}{2 \cos x} - \sin x \cos^2 x$$

$$\frac{\sin 2x}{2 \cos x} = \frac{2 \sin x \cos x}{2 \cos x} = \sin x$$

$$\sin x - \sin x \cos^2 x \quad \bullet^1 \checkmark$$

$$\sin x(1 - \cos^2 x) \quad \bullet^2 \checkmark$$

Question			Generic scheme	Illustrative scheme	Max mark
11.	(b)		<ul style="list-style-type: none"> •⁴ know to differentiate $\sin^3 x$ •⁵ start to differentiate •⁶ complete differentiation 	<ul style="list-style-type: none"> •⁴ $\frac{d}{dx}(\sin^3 x)$ •⁵ $3\sin^2 x \dots$ •⁶ $\dots \times \cos x$ 	3
Notes:					
Commonly Observed Responses:					

[END OF MARKING INSTRUCTIONS]